

minehen Aug 1869 Stat. Naturae Curios. vol 1X -10 (1752) append. pr 172 - 208 - tal 10-13 I. Phil. Bregnii descriptio fruction Cerci America.
mejoris articulati, flore maximo\_ norta se apericula suavissiume obsem spirante (Cer. grandiflonus) The plant often flowered, but never product fuit Sometim, 2-4 flowery opening on the Jane evining some seen in centril aftern open a July 27 1748, brought pail sipe 11 months afterwards. \_ pris 2 inches long, 1 - Samter spellowish flesh rolow, felenseleted, on each televele a bunk of world, hair - nets of flow tedhen hag -Juliof Black shiring seed in Milich prepsom of the already generating in the prop Follows a long descriptorin by Christ. Jar. Trew Jute Muraben Lan. 17.52 with many observations of figures. of Cercus grandiflers - others. He shows efigure, t. XI. f 2 That the acrice wit som from the grovers 503 also i the funite are bruncher: XII.f. 4 ad pry: 196 copia filorum ramosomme globuli, [ovila] onustone

copyright reserved cm


MISSOURI BOTANICAL GARDEN

At faith states that what he calls stamina serundiordinis ovivi ex Tubi lateribus ita ut illa ad unum pollirem supra ovarium t. XI. J. 10 E.B., et duos pollères infor conlle segmente, och distingen potævint stamme primi ordinis [ The Corona) al corolle segmentorum ortum dl. totan taben humore viscoso melles

1 ratere -

p. 197 muguan ex selvo, sumper ex acrie costre provinit flos et quiden nomise profime super quendan spinarme ferriculan, ali quoque rami Colyris segmenton- numer 1 58-64 corollae vero altra 28 Styli-findrice 17 nune plure, semel 23

MISSOURI cm copyright reserved GARDEN




### M. W. ALEXANDER,

UTH-EAST CORNER OF FOURTH AND MARKET STREETS, ST. LOUIS. ST. LOUIS.

MISSOURI BOTANICAL GARDEN GEORGE ENGELMANN PAPERS






# M. W. ALEXANDER.

THEAST CORNER OF FOURTH AND MARKET STREETS, ST. LOUIS.

MISSOURI BOTANICAL GARDEN GEORGE ENGELMANN PAPERS








# SOUTH-EAST CORNER OF FOURTH AND MARKET STREETS, ST. LOTTIS.


MISSOURI BOTANICAL GARDEN GEORGE ENGELMANN PAPERS











### M. W. ALEXANDER.

UTII-EAST CORNER OF FOURTH AND MARKET STREETS, ST. LOUIS.

MISSOURI BOTANICAL GARDERS GEORGE ENGELMANN PAPERS



Cerus grandiflomes fix vaised by the Cite Mr Griever Stanis 1839 1836 magnifi sorte of engy forced with short wooll, long hair a gometitus oftens yellowith Thinks 12 mora are the most detet 7 × 12 rows of sales on fit, crossing each other; about (12 armagem x); or 31 beller member of soules 130-150. tule of flower persot to frit scarlet contracts above interior pule greenish 2000 1,1-1,2 lin log bleck smooth shining with irragular & shallow ingeressions or dots MISSOURI BOTANICAL copyright reserved cm GARDEN

of the Chamiter of the vegetather?

of South water Texas

Frea of the M. S. shows of we may naturally ax feet several Different regions of vigetation product or influenced by Lititle, longitur conformation of ruspind, geologiste characte of the soil, and perhaps others faceres not fo verdig appreniable. The Hogetation of the immense territory of the U.S. bears a people en Character Distinguishing it from other regions of Ogalatin This character on it in the prevalence of some families and more especially of the fin the source of Compositue, and in the great number other, only in this of Speries of many genera, known part of the world 59. Magnolinea) ni other countries only by a glan representative or known out among the footil remains of the form coal (tation) formation. Juch chesnuts or Buckeyes the bisches, Walnut, & Histories, the mysolist & the granding. The goodshies and Succes grown (Liquidade).

are Vorly for among & formils, in the Lesbores plants examplifying Their GEORGE ENGELMANN PAPERS chaute are the Atin, Tolishing or, Kelsathen, Applepris st. Det as we may naturally infers their lage territory shows Different regions of vegetation, produced on inflerence by directe information of the Jungeon geologis-el character of the sail and purhaps other causes not 10 readily appreviable. I well Different are at are the north. eastern Affighings I thing slope to

the dellation - the south of but of left many & this easter the Trapind registate floor Floored tothe occasi MISSOURI

BOTANICAL copyright reserved cm GARDEN



copyright reserved

BOTANICAL GARDEN


Cereires significables duk Joebe C, Jand 25 5 1860 12 legt of flower 11 inhay ovarg globore jetak dånte, Blet 130 pulnie. in 1/34 arranged, techennelt of triangle weathers tule alex & into les le Celons frink this lefound widens to weary 2 in the with numerous 8000/00 hiangelan scales bearing wood and brittles - Hail 19 julium free spread sepel, all 50-60 4" 2 juliuhe & long line shorte 3" redrik, oulg 2 liver wide 26 lette, al 25 white of and and and to constitute 3 he with by 15-16 Cine Grows represents field, seminent, form a capel al 1 Le melen with The case'd neket fellow nertaniferon 1/2 in they the for 4 recher innumenable stances, naket space 1/4 inch et the corona of numeron stamen, alex 3 inder long stiffe con inher by whilif of eque thicken the stiguets creek petalon, ante, timen authen 1/4 lin les, Caren, emany, et Gove of The places ? Carche, To fall in separate pakes MISSOURI BOTANICAL cm copyright reserved GARDEN


Cereus nyetstallen at E danders duly 10'1863. Longth of own 1" Thicks " Lepal, topit, & petals pull skruk ut men at probably som 11" Eaght, of flow 10" Ovang globose, with abl 110-120 pulorlli 20 annuged - He 10 at 16 spirals most conspioner ovang strongly tuberralated, arcola on each Turente bearing a semilerete abolato scale Common, 1 enperous 2 likes long (resultes lenver of Operation, band in its axil white ship this some (4-6) slift white spines

or curly rouse white hair The 5" log with all 60 = 80 septs, in 10 amount 4 & 6 spring most conspicuous; the 16 verticely plainly mark of 16 sibs, sept All the factor of the flatter oblinseilate, and bearing in their axil, tometime i for white opines or long course carry bristles of I will long spread & recommend peterion sepel, zellow mend medlik och s line sabelet 50-00 i number, convergonés flat or sette some inside - 4-6 standet ongs bronde, estite from transitor to factal petal, estite oblacerole - spetalet, and 21-30 - number, fring a kir of som glober or caly x shaped rowlla. Alaka space of tale 1/2 inch, stame to some 3 ing apper make space also 1/4 inches - 88 stames - corona, 2/2-2/4 the Con- stige anthe, Ceniar 2/2 likes Cour (not 1/4 ason finkpage!) stigm. 12 linnen and MISSOURI

BOTANICAL copyright reserved cm GARDEN

estite aula Corus nyetirallus Hort Jouen Sept 1862 very vigorous. At joints oblinely quachy Lar almost rybudst-, young ony 4 winged, angles vompressed. brit sarely 5 augles. 1 7 2 3






20 Cin Cercus nyotipally? greaves Mov. 1858 4,048 fuit globone Monis 2 jacker L'aut anth resi vore rod with about 120/30 ones more appropries 1005 ell i 13/ annyt. all spring; 5- 8 yellowit Thomas 2-5 Chang · wile long, and long septs spran lower arcolae iz ship bu stiff outy hair. mully with 1 both mulle pruis a more approxity uppe erus mene rass of st Distant en ents flesky smles on the ter med purbly upp part of the fruit megore, or with slightly suloute from the protuberon of the app publity top daply umblink. Thin red pun'range. white pulp. Sert. 2 /4 black sends spanely Dith but MISSOURI BOTANICAL

copyright reserved

GARDEN



Vollen 17-20 Cin -0.041 -0,048 lie Namt The lose or south vestoles 22 31 1065 Cer, nyrhrellen at Shaws How 12 rache, log. Tube ? long with so ongo somes i 73, iaan depel 58 - 4 jahrtog, 3 l. wide, a far intermedition - Lyds spray 10-102 26 petet, langed one, 15-18 l. wide sprendje 4-42 inder long. yellow neitan form maked space - bottom of upper nek & spare between these of stamen a O Corona of Jann Leagth -Ile 1 100 stances ~ Corona 14 Stogmete 1263 2lg 5- 1860 445" 23/4 3/2-4/2 MISSOURI BOTANICAL copyright reserved

GARDEN

cm

